Evaluación de la estructura factorial de la prueba MAC-R (Cuestionario de Cogniciones Anoréxicas de Mizes) en población femenina adulta y adolescente de la Ciudad de Buenos Aires

Assessing the factorial structure of the MAC-R test (Mizes Anorectic Cognitions Questionnaire) in adolescent and adult female population of the City of Buenos Aires

Vanesa Carina Góngora[1]

RESUMEN

Este trabajo examina la estructura factorial de la prueba MAC-R que evalúa cogniciones específicas de los trastornos alimentarios (TCA) en población femenina adulta y adolescente de la ciudad de Buenos Aires. Se utilizaron tres grupos de mujeres: 447 adultas, 432 adolescentes y 95 pacientes con TCA. Se emplearon la escala MAC-R y la escala ICA de síntomas alimentarios. Se realizó un análisis factorial exploratorio por componentes principales con solución de 3 factores y de 2 factores y rotación Varimax. Se halló una solución de 15 ítems en 2 factores que explicaba 46.51 % de la varianza con coeficientes alfas de .83 y .77. El análisis factorial confirmatorio mostró que el modelo de dos factores se adecua mejor a los datos. La misma estructura se replicó en el grupo de adolescentes. La versión de 15 ítems en dos subescalas, Autocontrol alimentario y Aprobación, ha mostrado buenas evidencias de validez y confiabilidad.

Palabras clave: factorial – escala – cogniciones – validez - alimentación.

1 Dra. en Psicología, Investigadora del CONICET, docente de la Facultad de Psicología de la Universidad de Buenos Aires. E-mail: vgongora@psi.uba.ar

ABSTRACT

This study examines the factor structure of the MAC-R test that assesses specific cognitions of eating disorders (ED) in adolescent and adult female population of the city of Buenos Aires. Three female groups were used: 447 adults, 432 adolescents and 95 ED patients. The MAC-R scale and the ICA eating symptoms scale were administrated. An exploratory factor analysis by principal components with 3 factors and 2 factors solutions and Varimax rotation was performed. A two-factor solution was found containing 15 items, which explained 46.51% of variance. Alpha coefficients were .83 and .77. The confirmatory factor analysis showed that the two-factor model best fitted the data. The same factorial structure was replicated in the adolescent group. The 15 item version of the MAC-R with two subscales, Self-control and eating, and Approval, has shown good evidence of validity and reliability.

Keywords: factorial - scale - cognitions - validity – eating.